



A STUDY OF BRANCHING PATTERN OF EXTERNAL CAROTID ARTERY IN HUMAN CADAVERS

Anatomy

Dr. Sweta Rani	Junior Resident, Deptt of Anatomy DMCH, Darbhanga Laheriasarai
Dr. Seema Tabassum	Associate Prof. Deptt of Anatomy DMCH, Darbhanga Laheriasarai
Dr. Vandana Varsha*	Junior Resident, Deptt of Anatomy DMCH, Darbhanga Laheriasarai. *Corresponding Author
Dr. G.S.Jha	Assistant Prof. Deptt of Anatomy DMCH, Darbhanga Laheriasarai
Dr. S.K. Karn	Associate Prof and Head, Deptt of Anatomy DMCH, Darbhanga Laheriasarai

ABSTRACT

BACKGROUND: To study in the branching pattern of the external carotid artery, being the principal artery nourishing almost the head and neck (external to skull) and report any deflections from the normal branching pattern described in the text books of Anatomy. **MATERIALS, METHODS AND RESULTS:** Total study of 30 cadavers at DMCH, Laheriasarai, bihar (24 males and 6 females), we have found variations in the various branches and reported (figures).

KEYWORDS

External Carotid Artery (ECA), Common Carotid Artery (CCA), Superior Thyroid Artery, Ascending Pharyngeal Artery, Lingual Artery, Facial Artery, Occipital Artery

INTRODUCTION

The external carotid artery is a major artery of the head and neck. It arises from the common carotid artery when it splits into the external and internal artery. The external carotid artery supplies blood to the face and neck. The external carotid artery begins at the upper border of the thyroid cartilage and curves, passing forward and upward. And then including backward to the space behind the neck of the mandible, where it divides into the superficial temporal and maxillary artery within the parotid gland. A detailed knowledge of its branching pattern is of considerable significance in planning head and neck surgery.^[1] This also necessitates accurate interpretation of radiological images and proper planning of the surgical steps taken during the exploration of head and neck such as carotid endarterectomy, aneurysm repair, carotid angiogram and radical dissection of the lymph nodes.^[2,3,4] The proper identification of the branches of the external carotid artery is imperative.^[5,6] Branches of ECA are key landmarks for adequate dissection for exposure and appropriate placement of cross clamps on the carotid arteries or catheterization.^[7] The patterns of branching of ECA show ethnic differences.^[8] High variability of the carotid arterial system in the Kenyan population has been reported.^[9,10,11]

MATERIALS AND METHODS AND RESULTS


Study Site: Total 30 cadavers (24 males and 6 females) ages ranging from 35-50 years were dissected at the Deptt of Anatomy dissection hall, Darbhanga Medical College and hospital Laheriasarai bihar. One specimen showed a common stem arising from ECA trifurcated into ascending pharyngeal, lingual and facial arteries (Fig1). Ascending pharyngeal artery arose higher than lingual artery on one specimen and it branched from occipital artery as slender branch in one specimen (Fig.2a and 2b). In one specimen a short stem was given off from ECA which divided into two branches: one continued as the superior thyroid artery and the other trifurcated into ascending pharyngeal, lingual and facial arteries (Fig.3). In one specimen, lingual artery and facial artery formed a common linguofacial trunk. Facial artery in one specimen was given off at the level of the angle of the mandible much higher than its normal level of origin (Fig4a and 4b). Occipital artery branched off as a common trunk with posterior auricular artery in two specimens (Fig.5a). In another specimen, it arose as an independent branch below the linguofacial trunk (Fig.5b). In one specimen it was in between the superior thyroid and lingual arteries. All of the variations were observed on the left side.

Figures. 1.



A common trunk (*) trifurcating into lingual (2), ascending pharyngeal (3) and facial (4) arteries.

2.a



CCA – Common Carotid artery
 ECA – External Carotid artery
 ICA – Internal Carotid artery
 1. Superior thyroid artery
 2. Lingual artery
 3. Facial artery
 4. Continuation of External Carotid artery
 *Ascending Pharyngeal artery (Higher origin)



2.b

Ascending pharyngeal artery (1) stemming from occipital artery (2) 3.



A common trunk *giving off superior thyroid artery (1) before trifurca

DISCUSSION

The External Carotid Artery is one of the terminal branches of Common Carotid Artery usually given off at the level of the upper border of thyroid cartilage, corresponding to C3-C4 vertebral level.^[12] It is anteromedial to the internal carotid artery at its origin in the carotid triangle and ascends to the parotid gland and terminates into maxillary and superficial temporal arteries at the level of the neck of the mandible^[12]. It gives off superior thyroid, lingual, facial, ascending pharyngeal, occipital and posterior auricular arteries in the neck. In a radiological study in different age groups, it is reported that the origin of ECA can be anywhere between C2-C6 vertebral levels^[13]. In our present study the ECA was given off at the usual level in all the specimens. The variations observed in the present study are linguofacial trunk, trifurcation of a common trunk and higher and lower level of origins of some branches.

External Carotid Artery development is a complicated process of angiogenesis and remodelling which includes annexation and regression of vessels. The development of hyostapedial artery which links the neural crest arterial system to the ventral pharyngeal arterial system is an important event in the development of external carotid artery. Signals involved in annexation and regression are not always synchronized which results in various anatomical variations^[14]. A case report was found which showed an anomalous glandular branch arising

directly from the ECA, on the medial aspect, 1.2 cm above the bifurcation of CCA. It exclusively supplied the submandibular salivary gland^[15]. In a study done among 30 cadavers, a case was found to have glandular branches directly given off from the ECA to the parotid salivary gland. They also report about unusual origin and course of the right facial artery which arose from the ECA just above the angle of the mandible and passed directly on to the face crossing the posterior border of mandibular ramus and along the posterior pole of the superficial part of the submandibular salivary gland^[16].

CONCLUSION

The external carotid artery shows high frequency of variations comprising early bifurcation, trifurcation, quadrifurcation and short common stem. These variations may lead to inadvertent injury and cause confusion in interpretation of angiograms. Preoperative evaluation is recommended^[17]. In the present study we did not come across quadrifurcation. Bifurcation, trifurcation and higher and lower origin of some branches were observed.

For reconstruction purposes, the Facial Artery Musculo.Mucosal (FAMM) flap was introduced. But its use is limited by variations in the course of the facial artery. Therefore, application of knowledge of the precise course and branching pattern of the facial artery is required for construction of FAMM flap and its successful utilization^[18,19].

REFERENCES

- Hoffer SO, Poschn A, Smit X. 2005. The facial artery perforator flap for reconstruction of peroral defects. *Plast Reconstr Surg*; 115: 996 – 1003.
- Thwin SS, Soe MM, Myint M, Than M, Lwin S. 2010. Variations of the origin and branches of the external carotid artery in a human cadaver. *Singapore Med J*; 51: e40 – e42.
- Delic J, Savkovic A, Bajtarevic A, Isakovic E. 2010. Variations of ramification of external carotid artery – common trunks of collateral branches. *Period Biol*; 112: 117 – 119.
- Ozgun Z, Govsa F, Ozgur T. 2008a. Anatomic evaluation of the carotid artery bifurcation in cadavers: Implications for open and endovascular therapy. *Surg Rad Anat*; 30: 475 – 480.
- Sanjeev IK, Anita H, Ashwini M, Mahesh U, Rairam GB. 2010. Branching pattern of external carotid artery in human cadavers. *J Clin Diagn Res*; 4: 3128 – 3133.
- Panagouli E, Lolis E, Veneriatos D. 2011. A morphometric study concerning the branching points of the main arteries in humans: relationships and correlations. *An Anat*; 19: 86 – 99.
- Rao TR, Shetty P. 2011. Unusual Branching Pattern of External Carotid artery in a cadaver. *ChangGung Med J*; 34(6 suppl): 24 – 27.
- Sanjeev IK, Anita H, Ashwini M, Mahesh U, Rairam GB. 2010. Branching pattern of external carotid artery in human cadavers. *J Clin Diagn Res*; 4: 3128 – 3133.
- Anangwe D, Saidi H, Ogeng'o J, Awori KA. 2008. Anatomical variations of the carotid arteries in adult Kenyans. *E Afr Med J*; 85: 244 – 247.
- Ongeti KW, Ogeng'o JA. 2012. Variant origin of the superior thyroid artery in a Kenyan population. *Clin Anat*; 25: 198 – 202. [11]. Magoma G, Saidi H, Kaisha WO. 2012. Origin of thyroid arteries in a Kenyan population. *Annals Afr Surg*; 9: 50 – 54.
- Gray's anatomy: the anatomical basis of clinical practice. 40th ed. Standring S. eds. London: Elsevier Churchill Livingstone, 2008. [13]. Czervinski F. 1981. Variability of the course of external carotid artery and its rami in man in light of anatomical and radiological studies. *Folia Morphol*; 4, 449-453.
- Larsen W. Human Embryology. 2nd ed. New York: Churchill Livingstone, 1998: 191-195.
- Mamatha T, Rajalakshmi R, Latha VP, Gavishiddappa AH, Jiji P, Prameela MD. Anomalous branching pattern of the external carotid artery : a case report. *Rom J Morphol Embryol* 2010; 51(3): 593-595. (PMID: 20809046).
- Anu VR, Pai MM, Rajalakshmi R, Latha VP, Rajanigandha V, Costa DS. Clinically relevant variations of the carotid arterial system Introduction : Developmental anomalies. *Singapore Med J* 2007; 48(6): 566-569. (PMID: 17538758).
- Julius A. Ogeng'o, Musa K. Misiani, Poonamjeet Loyal, Kevin W. Ongeti, Jacob Gimongo, Martin I. Inyimili, Aclous K. Murunga. Variations in branching pattern of external carotid artery in a black Kenyan population. *Anatomy Journal of Africa*. 2015. Vol 4 (2): 584-590.
- Pribaz J, Stephens W, Crespo L, Gifford G. A new intraoral flap: facial artery musculomucosal (FAMM) flap. *Plast Reconstr Surg* 1992; 90(3): 421-429. (PMID: 1513887).
- Hima BA, Narsinga RB. Aberrant patterns of branching of external carotid artery. *Int J Basic Appl Med Sci* 2012; 2(2): 170-173.

4a.



Linguofacial trunk (2) 4b.



High origin of the facial artery* close to the angle of the mandible 5a.



A common occipito auricular trunk dividing in to occipital (1) and posterior auricular (2) arteries.

5b.



Low origin of occipital artery* below linguofacial trunk