



CARDIOVASCULAR RISK IN PATIENTS WITH SUBCLINICAL HYPOTHYROIDISM

Medicine

Dr. Anil Kumar Mehta*

M.D. (Medicine), DTM & H, Associate Professor, Department of Medicine, Darbhanga Medical College and Hospital, Laheriasarai, Bihar. *Corresponding Author

Dr. Debarshi Jana

Young Scientist (DST) Institute of Post-Graduate Medical Education and Research, A.J.C. Bose Road, Kolkata-700020, West Bengal, India.

ABSTRACT

Thyroid hormone regulate every organ system in our body. Metabolic parameters, including lipoprotein metabolism and cardiovascular risk factors like obesity, glucose intolerance or type 2 diabetes, dyslipidemia and hypertension are all influenced by serum TSH level in our body. Several reports have demonstrated a positive correlation between TSH levels and cardiovascular risk factors. C-reactive proteins is one of the markers of inflammation and also considered to be the predictor of cardiovascular disease and may be helpful in the assessment of future development of cardiovascular abnormalities in subclinical hypothyroidism.

Methods: Recently diagnosed 71 subclinical hypothyroidism patients were enrolled in the study and compared with 63 healthy individuals. Patients were divided into two groups on the basis of thyroid stimulating hormone (TSH) concentration. Thyroid profile (TSH, free thyroxine, triiodothyronine) and C-reactive protein were estimated by enzyme-linked immunosorbent assay. Total cholesterol (TC), triglycerides (TG), and high-density lipoprotein-cholesterol were estimated in the study population by a spectrophotometric method, and low-density lipoprotein-cholesterol (LDL-C) was measured by the Friedewald formula.

Results and Conclusion: Altered lipid profile, as well as higher concentrations of C-reactive protein, was observed in subclinical hypothyroidism patients as compared to controls. Higher concentration of TC, TG, and LDL was observed in subclinical hypothyroidism patients while the concentration of high-density lipoprotein was lower in patients with subclinical hypothyroidism. The elevated concentration of C-reactive protein was also observed in the patient group. Patients having TSH >10 μ U/ml have a higher risk of developing cardiovascular disease compared to patients having TSH <10 μ U/ml and increasing concentration of C-reactive protein along with LDL-C may to cardiovascular abnormality.

KEYWORDS

Subclinical hypothyroidism, C-reactive protein, Dyslipidemia, Cardiovascular risk.

INTRODUCTION

Subclinical hypothyroidism (SCH) is a common endocrine disorder presented by mildly elevated thyroid stimulating hormone (TSH) above the reference range along with a normal concentration of free thyroxine (FT4) and triiodothyronine (T3). It is an oscillated state between euthyroidism to hypothyroidism as it may lead to overt hypothyroidism. Thyroid hormones are known to play a significant role in cardiac hemodynamic because the deficient or excessive concentration of thyroid hormones may develop cardiac abnormalities. It is well-established that overt hypothyroidism is associated with cardiovascular symptoms through biochemical analysis. Dyslipidemia is a prominent feature of hypothyroidism established by various evidences, but not for the SCH due to inconsistent findings over the years.

The diversity of the result has raised the question that whether SCH is characterized by cardiovascular symptoms or not. C-reactive protein (CRP), an acute-phase reactant, is an effective and stable tool for the assessment of cardiovascular risk. It is an important marker of various inflammatory disorders like rheumatoid arthritis, myocardial infarction and various others etc. CRP is known to play an important role in the progression of atherosclerosis, a cardiovascular disease caused by the altered mechanism of low-density lipoprotein cholesterol (LDL-C). Therefore, the main objective of this study is an assessment of the role of CRP in SCH patients.

MATERIAL AND METHODS

This cross-sectional study was carried out in Department of Medicine, Darbhanga Medical College and Hospital, Laheriasarai, Bihar. Total 71 recently diagnosed SCH patients were enrolled for the study, in which 44 patients were having TSH 6.16- 10 μ U/ml and rest 27 patients were having TSH >10 μ U/ml termed as Cases-I and Cases-II respectively, compared with 63 healthy individuals as controls. People with a previous history or family history of thyroid diseases, taking thyroid medication, cardiovascular disease, smoking, menopausal women, pregnancy, inflammation, diabetes, hypertension, or alcoholics were excluded from this study. Age group criteria were kept from 21 to 45 years for the study population.

Body mass index (BMI) was calculated by means of height and weight of an individual. The weight of a person is divided by the square of the height. The weight and height of an individual were measured in kilogram (kg) and meter (m), respectively. The person having BMI

18.5 kg/m² to 24.99 kg/m² was considered a normal BMI. Thyroid profile (TSH, FT4 and T3) of SCH patients, as well as controls, was measured by enzyme-linked immunosorbent assay (ELISA). Patients with a concentration of TSH >6.2 μ U/ml along with a normal concentration of FT4 and T3 was considered to have SCH. ELISA kits was used for the investigation of thyroid profile estimation.

Total cholesterol (TC), triglycerides (TG), high-density lipoprotein cholesterol (HDL-C), and LDL-C were measured in the study group. TC, TG, and HDL-C were investigated by means of glucose oxidase (CHOD)/peroxidase (POD) method, GPO-PAP method and CHOD-POD/phosphotungstate method, respectively.

STATISTICAL ANALYSIS

All the variables (age, BMI, TSH, FT4, T3, lipid profile, and CRP) were expressed in mean \pm standard deviation (SD). One-way ANOVA was used for the differentiation of all the parameters between the various groups (Control group, Cases-I and Cases-II). A p<0.05 was considered statistically significant. IBM SPSS version 20 (Statistical package for social sciences) was used for the statistical analysis.

RESULT

This study shows a statistical difference among the various groups regarding the different variables. The age of all the participants was not significantly different statistically. Thyroid profile (TSH and T3) was significantly different among the groups except the FT4. BMI was significantly different among the groups and was highest in Cases-II and lowest in control groups (Table 1).

Lipid profile was significantly different among the groups. TC, TG, and LDL-C were higher in SCH groups compared to control group and highest in Cases-II. HDL-C was significantly different among the groups. The lowest concentration of HDL-C was observed in the second group of SCH patients. The concentration was also lower in Cases-I comparing to control group. The level of CRP was significantly different among the groups as its level was comparatively increasing in each group, from Cases-I to the Cases-II (Table 2).

Table 1 : Baseline characteristics among the various groups[‡]

Sl. No.	Parameters	Control (0.39-6.16 μ IU/ml)	Case I (6.16-10 μ IU/ml)	Case II (>10.0 μ IU/ml)	p-value
1	Age (years)	35.46 \pm 3.77	35.75 \pm 3.71	36.25 \pm 2.96	0.628
International Journal of Scientific Research					1

All the variables were expressed as mean±SD. ‡By analysis of variance, SD: Standard deviation, BMI: Body mass index, TSH: Thyroid stimulating hormone, FT4: Free thyroxine, T3: Triiodothyronine

Table 2 :Lipid profile and CRP among the various groups‡

	BMI (kg/m ²)	22.68±1.82	26.4±2.13	29.64±3.83	<0.01
2	TSH (mIU/ml)	2.75±0.78	8.20±0.82	14.86±2.95	<0.01
3	FT4 (ng/dl)	1.15±0.20	1.15±0.24	1.25±0.30	0.196
4	T3 (ng/dl)	1.08±0.28	0.95±0.18	0.93±0.15	<0.01

All the variables were expressed as mean±SD, ‡By analysis of variance, CPR: C-reactive protein, TC: Total cholesterol, TG: Triglycerides, HDL: High-density lipoprotein, LDL: Low-density lipoprote

DISCUSSION

This study shows that SCH patients are characterized by increased BMI, lipid profiles, and CRP level. Rotterdam Study specified that subclinical hypothyroidism is a strong predictor of myocardial infarction and risk of atherosclerosis. Elevated TSH is found to be associated with increased BMI in SCH patients. Srideviet *et al.* supported our study by signifying altered concentration of lipid profile as characterized by significantly increased level of TC, a higher concentration of TG and elevated LDL-C. Similar to this study Erdemet *et al.* observed that the concentration of HDL-C was significantly lower in SCH patients compared to controls. Study of Kvetnyet *et al.* also support this study that SCH might be a risk factor for the development of cardiovascular diseases due to increased level of TG and CRP. CRP, a plasma protein, predominantly synthesized in liver, emerged as an additional cardiovascular risk factor was found to be significantly higher in the SCH group compared to control group.

CRP has been reported to bind to oxidized LDL and actively participates in atherogenesis by direct influence on complement system, vascular cell activation, lipid accumulation and thrombosis. The level of CRP was significantly higher in SCH patients promising the approach for further development of cardiac risk, supported by Gaoet *et al.* study. Gupta *et al.* also observed the increased concentration of inflammatory markers in SCH patients. While contrary to it other studies defined that SCH has no relation with dyslipidemia. Luboshitzkyet *et al.* described that CRP does not play any role to contribute to the increased risk of cardiovascular disease in SCH patients. Toruneret *et al.* described that SCH is characterized by dyslipidemia, but this association does not depend on the range of TSH. This study proposed that SCH patients having TSH>10 µIU/ml have a higher risk of future development of cardiac abnormalities matched with Hernandez-Mijares and Marwaha study. In addition to this patients having SCH with TSH<10 µIU/ml also showed some indication of altered lipid fractions, comparing to control group supported by Regmi study.

CONCLUSION

The outcome of this study specifies that SCH patients are characterized by increased BMI, dyslipidemia, and a higher concentration of CRP. This significant increase in CRP level might be the cause of development of cardiac symptoms in the coming future. Cardiovascular abnormalities can be developed in the later stage of disease. Although there are remarkable changes in different variables in SCH patients having TSH >10 µIU/ml, Patients with SCH in which TSH range was <10 µIU/ml also has some alteration in different variables compared to normal healthy adults. Increased BMI, altered lipid fractions and abnormal CRP might be helpful in the early detection to diagnose cardiovascular risk in SCH patients. The sample size of this study was small, so the further studies with larger number of patients will be helpful to establish the fact.

REFERENCES

1. Abbas JM, Chakraborty J, Akanji AO, Doi SA. Hypothyroidism results in small dense LDL independent of IRS traits and hypertriglyceridemia. *Endocr J* 2008;55(2):381-9.
2. Calabrò P, Golia E, Yeh ET. Role of C-reactive protein in acute myocardial infarction and stroke: Possible therapeutic approaches. *Curr Pharm Biotechnol* 2012;13(1):4-16.
3. Chang MK, Binder CJ, Torzewski M, Witztum JL. C-reactive protein binds to both oxidized LDL and apoptotic cells through recognition of a common ligand: Phosphorylcholine of oxidized phospholipids. *Proc Natl Acad Sci U S A* 2002;99:13043-8.
4. Christ-Crain M, Meier C, Guglielmetti M, Huber PR, Riesen W, Staub JJ, *et al.* Elevated C-reactive protein and homocysteine values: Cardiovascular risk factors in hypothyroidism? A cross-sectional and a double-blind, placebo-controlled trial. *Atherosclerosis* 2003;166(2):379-86.
5. Erdem TY, Ercan M, Ugurlu S, Balci H, Acbay O, Gundogdu S. Plasma viscosity, an early cardiovascular risk factor in women with subclinical hypothyroidism. *Clin Hemorheol Microcirc* 2008;38(4):219-25.
6. Erhardt JG, Estes JE, Pfeiffer CM, Biesalski HK, Craft NE. Combined measurement of

- ferritin, soluble transferrin receptor, retinol binding protein, and C-reactive protein by an inexpensive, sensitive, and simple sandwich enzyme-linked immunosorbent assay technique. *J Nutr* 2004;134(11):3127-32.
7. Fiarresga AJ, Feliciano J, Fernandes R, Martins A, Pelicano N, Timóteo AT, *et al.* Relationship between coronary disease and subclinical hypothyroidism: An angiographic study. *Rev Port Cardiol* 2009;28:535-43.
8. Gan SD, Patel KR. Enzyme immunoassay and enzyme-linked immunosorbent assay. *J Invest Dermatol* 2013;133(9):e12.
9. Gao CX, Yang B, Guo Q, Wei LH, Tian LM. High thyroid-stimulating hormone level is associated with the risk of developing atherosclerosis in subclinical hypothyroidism. *Horm Metab Res* 2015;47(3):220-4.
10. Garrow JS, Webster J. Quetelet's index (W/H²) as a measure of fitness. *Int J Obes* 1985;9(2):147-53.
11. Gupta G, Sharma P, Kumar P, Itagappa M. Study on subclinical hypothyroidism and its association with various inflammatory markers. *J Clin Diagn Res* 2015;9(11):BC04-6.
12. Hak AE, Pols HA, Visser TJ, Drexhage HA, Hofman A, Witteman JC. Subclinical hypothyroidism is an independent risk factor for atherosclerosis and myocardial infarction in elderly women: The Rotterdam Study. *Ann Intern Med* 2000;132(4):270-8.
13. Hernández-Mijares A, Jover A, Bellod L, Bañuls C, Solá E, Veses S, *et al.* Relation between lipoprotein subfractions and TSH levels in the cardiovascular risk among women with subclinical hypothyroidism. *Clin Endocrinol (Oxf)* 2013;78(5):777-82.
14. Hueston WJ, King DE, Geesey ME. Serum biomarkers for cardiovascular inflammation in subclinical hypothyroidism. *Clin Endocrinol (Oxf)* 2005;63(5):582-7.
15. Kvetny J, Heldgaard PE, Bladbjerg EM, Gram J. Subclinical hypothyroidism is associated with a low-grade inflammation, increased triglyceride levels and predicts cardiovascular disease in males below 50 years. *Clin Endocrinol (Oxf)* 2004;61(12):232-8.
16. Luboshitzky R, Herer P. Cardiovascular risk factors in middle-aged women with subclinical hypothyroidism. *NeuroEndocrinol Lett* 2004;25(4):262-6.
17. Marwaha RK, Tandon N, Garg MK, Kanwar R, Sastry A, Narang A, *et al.* Dyslipidemia in subclinical hypothyroidism in an Indian population. *Clin Biochem* 2011;44(14-15):1214-7.
18. Otterness IG. The value of C-reactive protein measurement in rheumatoid arthritis. *Semin Arthritis Rheum* 1994;24(2):91-104.
19. Pearce EN. Hypothyroidism and dyslipidemia: Modern concepts and approaches. *Curr Cardiol Rep* 2004;6(6):451-6.
20. Regmi A, Shah B, Rai BR, Pandeya A. Serum lipid profile in patients with thyroid disorders in central Nepal. *Nepal Med Coll J* 2010;12(4):253-6.
21. Ridker PM. Cardiology Patient Page. C-reactive protein: A simple test to help predict risk of heart attack and stroke. *Circulation* 2003;108(12):e81-5.
22. Rifai N, Warnick GR, Remaley AT. Lipids, lipoproteins, apolipoproteins, and other cardiovascular risk factors. In: Burtis CA, Ashwood ER, Bruns DE, editors. *Teitz Fundamentals of Clinical Chemistry*. Pennsylvania: Saunders An Imprint of Elsevier Inc.; 2010. p. 422-4.
23. Shekhar R, Chowdhary NV, Das MC, Vidya D, Prabodh S. Prevalence of subclinical hypothyroidism in coastal Andhra Pradesh. *Biomed Res* 2011;22(4):471-4.
24. Singh SK, Suresh MV, Voleti B, Agrawal A. The connection between C-reactive protein and atherosclerosis. *Ann Med* 2008;40(2):110-20.
25. Singh U, Dasu MR, Yancey PG, Afify A, Devaraj S, Jialal I. Human C-reactive protein promotes oxidized low density lipoprotein uptake and matrix metalloproteinase-9 release in Wistar rats. *J Lipid Res* 2008;49:1015-23.
26. Solanki A, Bansal S, Jindal S, Saxena V, Shukla US. Relationship of serum thyroid stimulating hormone with body mass index in healthy adults. *Indian J Endocrinol Metab* 2013;17(Suppl 1):S167-9.
27. Surks MI, Ortiz E, Daniels GH, Sawin CT, Col NF, Cobin RH, *et al.* Subclinical thyroid disease: Scientific review and guidelines for diagnosis and management. *JAMA* 2004;291(2):228-38.
28. Toruner F, Altinova AE, Karakoc A, Yetkin I, Ayvaz G, Cakir N, *et al.* Risk factors for cardiovascular disease in patients with subclinical hypothyroidism. *Adv Ther* 2008;25:430-7.
29. Tuzcu A, Bahceci M, Gokalp D, Tuzun Y, Gunes K. Subclinical hypothyroidism may be associated with elevated high-sensitive c-reactive protein (low grade inflammation) and fasting hyperinsulinemia. *Endocr J* 2005;52(1):89-94.
30. Udapa SV, Manjrekar PA, Udapa VA, Vivian D, Altered fructosamine and lipid fractions in subclinical hypothyroidism. *J Clin Diagn Res* 2013;7(1):18-22.
31. Vargas-Uricoechea H, Sierra-Torres CH. Thyroid hormones and the heart. *Horm Mol Biol Clin Invest* 2014;18(1):15-26.
32. Wang CY, Chang TC, Chen MF. Associations between subclinical thyroid disease and metabolic syndrome. *Endocr J* 2012;59(10):911-7.
33. Yeh ET. A new perspective on the biology of C-reactive protein. *Circ Res* 2005;97(7):609-11.